



# TROOPMAX

## HOW DOES IT WORK?

**TROOPMAX** is a bacterial larvicide with a high target specificity, its impact on non-target organisms and the environment is negligible.

## COMPOSITION

TroopMax 0.8* 10 <sup>8</sup> CFU/g	(%) w/w
<i>Metarhizium anisopliae</i> (spores)	2%, 1 X 10 <sup>8</sup> CFU/g
<i>Beauveria bassiana</i> (spores)	2%, 1 X 10 <sup>8</sup> CFU/g
<i>Bacillus thuringiensis var Kurstaki</i> (spores)	2%, 1 X 10 <sup>8</sup> CFU/g
<i>Verticillium lecanii</i> (spores)	2%, 1 X 10 <sup>8</sup> CFU/g
Carrier i (Dextrose)	q.s.

## DOSAGE & APPLICATION

5g per liter of water

Recommended dosage is for guideline purpose only. More effective application rates may exist depending on specific circumstances.

## TARGET SPECIES

Hemiptera, Diptera, Coleoptera, Lepidoptera, Orthoptera and Hymenoptera etc. Spittlebugs (Cercopidae), pine moth *Dendrolimus* spp., corn borers *Ostrinia nubilalis*, *Ostrinia furnacalis*; *Locustana pardalina*, *Locusta migratoria* *Chortoicetes terminifera*, *L. migratoria* *Anacridium melanorhodon*, *Hieroglyphus daganensis* *Schistocerca gregaria*, *Helicoverpa armigera*, *Spodoptera litura*, *Aphis craccivora*, *A. gossypii*, *Rhopalosiphum maidis*, *Brevocoryne brassicae*, *Myzus persicae* and *Bemisia tabaci* etc.

## USAGE AND STORAGE

Troopmax is compatible with all natural fertilizers. Maintain the product out of direct sunlight and store in a dark and cool place between 5–25°C (40–77°F). Do not refrigerate or freeze. Keep container tightly sealed after use. Keep away from children and pets. Do not inhale or ingest.