



# BUDMAX



3 EASY STEPS TO GROW KING SIZED B U D S

BUDMAX

Inoculate the soil with this myco-active formula, to gain an early advantage in your crop cycle. Longer roots means a greater rhizosphere. Hence, ensuring that your plant will be able to maximize it's nutrients uptake, resulting in strong and healthy growth and bloom cycle.

#### Dosage and method of application

ROOTX should ideally be applied to the roots or rootball during transplanting

Transplanting container grown plants, potting on etc.

Tease plant roots from the root ball if necessary and dust TNC ROOTX directly onto the roots, add the remainder to the planting hole.

Seedlings / Saplings Add ROOTX to the planting hole: 1 tsp (5g) per plant.

Bare Root Plants / Whips Apply directly to the roots allow any excess to fall into the planting hole. Cuttings
Dip the cutting in ROOTX before planting.





Consists of bioactive humic and fulvic substances of vermicompost origin. It consists of cytokinins, auxins, betaines and gibberellins that are derived from seaweed fermentation. It consists of biologically derived N,P,K and trace elements from vermi compost and seaweed which aid in better root and shoot growth and supplement the plant with essential nutrients at critical stages of crop growth.

#### Dosage and method of application

GROWX should be mixed with water and applied as a soil inocolant to the planting soil during the vegetative stage.

Early growth - Mix 5ml (1 tsp) of GROWX to 1L of water. Apply to the planting soil once every 2 weeks.

Mature growth - Mix 5ml (1 tsp) of GROWX to each litre of water. Apply to the planting soil once every week.



## **BLOOM** X

BLOOMX Influences the microbial environment in a way that the constructive microorganisms become dominant. This creates an environment, in which the microorganisms play a positive role in plant growth, plant quality and soil fertility by using fermentation. Fermentative decomposition is stimulated and decomposition disappears, so that less energy is lost.

Soil in which constructive microorganisms dominate can produce optimal productivity levels, suppress illness and produce high quality products.

#### Dosage and method of application

Mix 3g (1/2 tsp) of BLOOMX with 1L of water. Apply to the planting soil every 2 weeks.







101, Blue Bell Building, Sitaram Compound, Crawford Market, Mumbai - 400001, India.



### **BUDMAX CANNABIS KIT**

#### **ROOTX**

**Soil Treatment** – Mix 4 kg per acre in 50 Kg of well decomposed Fym /compost / vermi compost/field soil and incorporate in the soil before sowing / transplant.

Broadcast the above mixture in standing crop 25-30 days after sowing.

#### **BLOOM X**

**Soil Treatment** – 40 ml per Acre. Mix 40 ml in 20 Litres of water before using in 1 acre of land.

#### **ATTENTION**

This specimen label is provided for general information only.

- This biological product may not yet be available or approved for sale or use in your area.
- It is your responsibility to follow all Federal, state/provincial, and local laws and regulations regarding the use of biologicals.
- · Before using any biological, be sure the intended use is approved in your state/province or locality.
- Your state/province or locality may require additional precautions and instructions for use of this product that are not included here.
- Indo Gulf Comapny does not guarantee the completeness or accuracy of this specimen label. The information found in this label may differ from the information found on the product label.
- You should not base any use of a similar product on the precautions,
- instructions for use or other information you find here.
- Always follow the precautions and instructions for use on the label of the biological you are using.

For application to seed in commercial seed treating equipment of for use in on-farm planting equipment.

Microbial seed inoculant for improving nutrient availability for increased yield potential.

NONPLANT FOOD INGREDIENT Not a fertilizer substitute

Warning: May form combustible dust concentrations in air [with packages greater than 25 kg].

Read the entire label before using this product. Use only according to label directions. Open package only when ready to use. Use before expiration date.

Apply this product only as specified on this label. Reference the specified application rate to determine the proper amount needed for seed. Additional talc and/or graphite may be applied if needed.

Dust particles may cause slight eye irritation. May cause respiratory irritation. Causes damage to lungs by inhalation through prolonged or repeated exposure.

This product may cause adverse effects to individuals who are repeatedly exposed to it. Eye or skin contact, inhalation, or ingestion should be avoided. Do not eat, drink or smoke when using this product. Wear protective gloves, protective clothing, and eye protection. Avoid breathing dust. Ensure adequate ventilation. Use only outdoors or in a well-venti lated area. Where exposure through inhalation may occur, use respiratory protective equipment. Wash thoroughly after handling. If inhaled: Remove person to fresh air and keep comfortable for breathing. Get medical advice/attention if you feel unwell.

The active ingredients in product are Bacillus amyloliquefaciens and Trichoderma virens spores. This product contains microorganisms that may have the potential to provoke sensitizing reactions; use appropriate personal protective equipment (PPE) to reduce exposure .Sensitized individuals should wear an approved respirator. In case of contact with skin or eyes, immediately flush exposed areas with plenty of water. Get medical attention if irritation occurs.

# Applicators and Other Handlers

Product contains live spores of the naturally occurring fungus Trichoderma virens and the naturally occurring bacteria Bacillus amyloliquefaciens, which may be damaged when exposed to high temperatures or contact with water for more than 4 hours. The live organisms contained in Product require specific storage conditions to ensure viability and product performance. To maintain product viability:

- Store unopened containers of Product in a dry cool place (less than 68"F or 20"C) away from sunlight and heat sources. Store locked up.
- Minimize temperature fluctuations.
- Avoid freeze/thaw cycles.
- Use before the expiration date. The expiration date is valid for unopened containers stored according to conditions listed above.
- Keep out of reach of children.

Unused Product and seed to which Product has been applied should be disposed of in accordance with applicable Federal, state/provincial, and local requirements.

# <u>Limited Warranty</u>

The Seller warrants that this product contains a minimum number of Bacillus amyloliquefaciens and Trichoderma virens colony forming units, as specified on this label. The Seller makes no other warranty expressed orimplied as to product viability or performance since product storage, use and growing conditions are beyond the Seller's control. Seller's guarantee is limited to the terms set out on the label and subject thereto. Buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.If this product does not perform as warranted above, and to the extent consistent with applicable law, customer's sole remedy for breach of warranty shall be replacement of the product or refund of the purchase price paid, at the option of Indo Gulf Company.

#### BIOTECH DIVISION OF INDO GULF COMPANY

101, Blue Bell Building, Sitaram Compound, Crawford Market, Mumbai 400001, India.

T:+91-22-23455354 | F:+91-22-23476536 E:indogulf@gmail.com | W:www.indogulfbioag.com

